























- Bahjat KS, Freitag NE, Brockstedt DG and Dubensky TW. Constitutive activation of the PrfA regulon enhances the potency of vaccines based on live-attenuated and killed but metabolically active *Listeria monocytogenes* strains. *Infect Immun* 2008; 76: 3742-3753.
- [16] Gedde MM, Higgins DE, Tilney LG, and Portnoy DA. Role of listeriolysin O in cell-to-cell spread of *Listeria monocytogenes*. *Infect Immun* 2000; 68: 999-1003.
- [17] Harding CV and Geuze HJ. Class II MHC molecules are present in macrophage lysosomes and phagolysosomes that function in the phagocytic processing of *Listeria monocytogenes* for presentation to T cells. *J Cell Biol* 1992; 119: 531-542.
- [18] Carrasco-Marin E, Shimizu J, Kanagawa O and Unanue ER. The class II MHC I-Ag7 molecules from non-obese diabetic mice are poor peptide binders. *J Immunol* 1996; 156: 450-458.
- [19] Forestier C, Deleuil F, Lapaque N, Moreno E and Gorvel JP. *Brucella abortus* lipopolysaccharide in murine peritoneal macrophages acts as a down-regulator of T cell activation. *J Immunol* 2000; 165: 5202-10.
- [20] Bruder D, Nussbaum AK, Gakamsky DM, Schirle M, Stevanovic S, Singh-Jasuja H, Darji A, Chakraborty T, Schild H, Pecht I and Weiss S. Multiple synergizing factors contribute to the strength of the CD8+ T cell response against listeriolysin O. *Inter Immunol* 2005; 18: 89-100.
- [21] Zenewicz LA and Shen H. Innate and adaptive immune response to *Listeria monocytogenes*: a short overview. *Microbes Infect* 2007; 9: 1208-1215.
- [22] Corr SC and O'Neill LA. *Listeria monocytogenes* infection in the face of innate immunity. *Cellular Microbiol* 2009; 11: 703-709.
- [23] Cervantes J, Nagata T, Uchijima M, Shibata K and Koide Y. Intracytosolic *Listeria monocytogenes* induces cell death through caspase-1 activation in murine macrophages. *Cell Microbiol* 2008; 10: 41-52.
- [24] Schnitzer JK, Berzel S, Fajardo-Moser M, Remer KA and Moll H. Fragments of antigen-loaded dendritic cells (DC) and DC-derived exosomes induce protective immunity against *Leishmania major*. *Vaccine* 2010; 28: 5785-5793.
- [25] Beauvillain C, Ruiz S, Guiton R, Bout D and Dimier-Poisson I. A vaccine based on exosomes secreted by a dendritic cell line confers protection against *T. gondii* infection in syngeneic and allogeneic mice. *Microbes Infect* 2007; 9: 1614-162.
- [26] Beauvillain C, Juste MO, Dion S, Pierre J and Dimier-Poisson I. Exosomes are an effective vaccine against congenital toxoplasmosis in mice. *Vaccine* 2009; 27: 1750-1757.