Product Info. Antibody Ver. 03/05/2012



Viale Piceno 137/f 61032 Fano PU (IT) Telephone + 39 (0)721830605 FAX +39 (0)721837154 e-mail:info@diatheva.com www.diatheva.com



Mouse Anti HIV-I p24 (clone 2)

ANT0068 50µg

Description	HIV-1 Gag p24 antigen is a capsid protein that constitutes the core of HIV-I. P24 protein is detectable starting from the early stage of AIDS virus infection, for this reason it is an interesting marker for disease progression in infected patients. Monoclonal anti p24 antibody clone 2 is designed for the detection of p24 core antigen in human serum or plasma from infected individuals as well as in cell culture supernatant.
Product type	Monoclonal antibody. (Hybridoma provided from King's College London Business Ltd. Dr Michael Jorgensen)
Immunogen	Bacterially expressed, hexahistidine amino-terminal tagged HIV-1 p24 gag recombinant protein (clade B, HXB-3 isolate).
Source	Mouse monoclonal IgG _{1,k}
Reacts with	HIV-1 p24 protein
Specificity	NA Epitope map not available.
Tested applications	ELISA, Western Blotting, Immunofluorescence
Recommended dilutions	Recommended starting dilutions can vary lot-to-lot. Consult the product information label in the package for lot specific values.
Recommended dilutions	
Recommended dilutions	Consult the product information label in the package for lot specific values. Note: When using any primary antibody or fluorescence-labelled secondary antibody for the first time, titrate out the antibody to determine which dilution allows the strongest
Recommended dilutions Purity	Consult the product information label in the package for lot specific values. Note: When using any primary antibody or fluorescence-labelled secondary antibody for the first time, titrate out the antibody to determine which dilution allows the strongest specific signal with the lowest background for your sample.
	Consult the product information label in the package for lot specific values. Note: When using any primary antibody or fluorescence-labelled secondary antibody for the first time, titrate out the antibody to determine which dilution allows the strongest specific signal with the lowest background for your sample. <i>For untested applications or species please refer to the <u>S.M.A.K.</u> program.</i> Antibodies are purified from supernatants of hybridoma cell cultures by affinity
Purity	Consult the product information label in the package for lot specific values. Note: When using any primary antibody or fluorescence-labelled secondary antibody for the first time, titrate out the antibody to determine which dilution allows the strongest specific signal with the lowest background for your sample. <i>For untested applications or species please refer to the <u>S.M.A.K.</u> program. Antibodies are purified from supernatants of hybridoma cell cultures by affinity chromatography.</i>